

Doset Hibi rings with an application to invariant theory

Mitsuhiro MIYAZAKI

Dept. Math., Kyoto University of Education,
Fushimi-ku, Kyoto, 612-8522 Japan
E-mail: g53448@kyokyo-u.ac.jp

Abstract

We define the concept of a doset Hibi ring and a generalized doset Hibi ring which are subrings of a Hibi ring and are normal affine semi-groupings. We apply the theory of (generalized) doset Hibi rings to analyze the rings of absolute orthogonal invariants and special orthogonal invariants and show that these rings are normal and Cohen-Macaulay and has rational singularities if the characteristic of the base field is zero and is F-rational otherwise. We also state a criterion of Gorenstein property of these rings.

Keywords: Hibi ring, sagbi, Doset, ring of invariants, Schubert subvariety

1 Introduction

Grassmannians and their Schubert subvarieties are fascinating objects and attract many mathematicians. Let $G_{m,n}$ be the Grassmannian consisting of all m -dimensional vector subspaces of an n -dimensional vector space V over a field K . Then the homogeneous coordinate ring of $G_{m,n}$ is $K[\Gamma(X)]$, where $X = (X_{ij})$ is the $m \times n$ matrix of indeterminates and $\Gamma(X)$ is the set of maximal minors of X .

For a Schubert subvariety of $G_{m,n}$, there correspond integers $b_1, \dots, b_m$ with $1 \leq b_1 < \dots < b_m \leq n$ and the universal $m \times n$ matrix Z with entries in a K -algebra S with the condition

$$I_i(Z_{\leq b_i-1}) = (0) \quad \text{for } i = 1, \dots, m,$$

where $Z_{\leq b_i-1}$ stands for the $m \times (b_i - 1)$ matrix consisting of the first $b_i - 1$ columns of Z and $I_i(Z_{\leq b_i-1})$ stands for the ideal of S generated by the i -minors of $Z_{\leq b_i-1}$. And the homogeneous coordinate ring of the Schubert subvariety of $G_{m,n}$ is $K[\Gamma(Z)]$, where $\Gamma(Z)$ is the set of maximal minors of Z .

On the other hand, because of the universal property of Z , any subgroup of $\text{GL}(m, K)$ acts on $K[Z]$. In this paper, we study the rings of absolute

$O(m)$ and $SO(m)$ -invariants (see Definition 3.1), denoted as $K[Z]^{O(m,-)}$ and $K[Z]^{SO(m,-)}$ respectively, of $K[Z]$.

We now state the contents of this paper. We first establish notation, recall known facts and state basic facts in section 2 in order to simplify the proofs and clarify the structure of the theory of the main part of this paper, sections 3, 4 and 5.

In section 3, we show that the ring of absolute $O(m)$ and $SO(m)$ -invariants are precisely the expected ones, that is, $K[Z]^{O(m,-)} = K[Z^T Z]$ and $K[Z]^{SO(m,-)} = K[Z^T Z, \Gamma(Z)]$. See (2.5) for notation.

In section 4, we introduce the concept of doset Hibi rings and their generalization. They are subrings of a Hibi ring and are normal affine semigroups, therefore are Cohen-Macaulay. We apply the theory of doset Hibi rings and generalized doset Hibi rings to investigate the property of $K[Z]^{O(m,-)}$ (or $K[Z]^{SO(m,-)}$ resp.) by showing it is isomorphic to a subring of a polynomial ring over K , whose initial algebra is a doset Hibi ring (or a generalized doset Hibi ring resp.) in an appropriate monomial order. In particular, $K[Z]^{O(m,-)}$ (or $K[Z]^{SO(m,-)}$ resp.) is a normal Cohen-Macaulay ring and has rational singularities if $\text{char} K = 0$ and is F-rational if $\text{char} K > 0$.

Finally in section 5, we state a criterion of Gorenstein property of a doset Hibi ring and a generalized doset Hibi ring. As an application, we state a criterion of Gorenstein property of $K[Z]^{O(m,-)}$ (or $K[Z]^{SO(m,-)}$ resp.), see Theorems 5.6 and 5.7. We note that the criterion of Gorenstein property of $K[Z]^{O(m,-)}$ is already obtained by Conca [Con]. Therefore, Theorem 5.6 recovers the result of Conca.

2 Preliminaries

In this section, we establish notation, recall known results and show some basic facts in order to prepare the main part of this paper, sections 3, 4 and 5.

2.1 Normal affine semigroups

(2.1) In this paper, all rings and algebras are commutative with identity element.

We denote by $\mathbf{N}$ the set of non-negative integers, by $\mathbf{Z}$ the set of integers, by $\mathbf{R}$ the set of real numbers and by $\mathbf{R}_{\geq 0}$ the set of non-negative real numbers.

Let K be a field and $X_1, \dots, X_r$ indeterminates, S a finitely generated additive submonoid of $\mathbf{N}^r$. We set $K[S] := K[X^{\mathbf{s}} \mid \mathbf{s} \in S]$ where $X^{\mathbf{s}} = X_1^{s_1} \cdots X_r^{s_r}$ for $\mathbf{s} = (s_1, \dots, s_r)$. Then

Theorem 2.1 ([Hoc]) (1) $K[S]$ is normal if and only if $S = \mathbf{Z}S \cap \mathbf{R}_{\geq 0}S$.

(2) If $K[S]$ is normal, then it is Cohen-Macaulay.

If $S = \mathbf{Z}S \cap \mathbf{R}_{\geq 0}S$, then by Farkas' theorem and the fundamental theorem of abelian groups, S is isomorphic to a toroidal monoid in the terminology of Stanley [Sta, p. 81]. So by [Sta, p. 82], we have the following

Theorem 2.2 If $K[S]$ is normal, then the canonical module of $K[S]$ is

$$\bigoplus_{\mathbf{s} \in \text{relint} \mathbf{R}_{\geq 0}S \cap S} KX^{\mathbf{s}}.$$

Stanley also stated, in the same article, a criterion of Gorenstein property for graded Cohen-Macaulay domains.

Theorem 2.3 ([Sta, 4.4 Theorem]) Let A be an $\mathbf{N}$ -graded Cohen-Macaulay domain with $A_0 = K$. Suppose $\dim A = d$. Then A is Gorenstein if and only if there is $\rho \in \mathbf{Z}$ such that $\text{Hilb}(A, 1/\lambda) = (-1)^d \lambda^\rho \text{Hilb}(A, \lambda)$, where Hilb stands for the Hilbert series.

As a corollary, we have

Corollary 2.4 Let A and B be $\mathbf{N}$ -graded Cohen-Macaulay domains with $A_0 = B_0 = K$. Suppose $\text{Hilb}(A, \lambda) = \text{Hilb}(B, \lambda)$. Then A is Gorenstein if and only if B is Gorenstein. In particular, if A is a graded subring of a polynomial ring with monomial order and the initial algebra in A of A is finitely generated over K and Cohen-Macaulay, then A is Gorenstein if and only if $\text{in } A$ is Gorenstein.

Note, by the flat deformation argument, we see that if A is an $\mathbf{N}$ -graded subalgebra of a polynomial ring with monomial order and $\text{in } A$ is Cohen-Macaulay, then A is Cohen-Macaulay.

2.2 Posets and lattices

(2.2) Let P be a finite partially ordered set (poset for short).

The length of a chain (totally ordered subset) X of P is $\#X - 1$, where $\#X$ is the cardinality of X .

The rank of P , denoted by $\text{rank}P$, is the maximum of the lengths of chains in P .

A poset is said to be pure if its all maximal chains have the same length.

A poset ideal of P is a subset I of P such that $x \in I$, $y \in P$ and $y \leq x$ imply $y \in I$.

For $x, y \in P$, y covers x , denoted by $x \triangleleft y$, means $x < y$ and there is no $z \in P$ such that $x < z < y$.

For $x, y \in P$ with $x \leq y$, we set $[x, y]_P := \{z \in P \mid x \leq z \leq y\}$.

A map $\nu: P \rightarrow \mathbf{N}$ is order reversing if $x \leq y$ in P implies $\nu(x) \geq \nu(y)$ and strictly order reversing if $x < y$ in P implies $\nu(x) > \nu(y)$.

The set of all order reversing maps from P to $\mathbf{N}$ is denoted by $\overline{\mathcal{T}}(P)$. The set of all strictly order reversing maps from P to $\mathbf{N} \setminus \{0\}$ is denoted by $\mathcal{T}(P)$.

For a family of indeterminates $\{T_x\}_{x \in P}$ indexed by P and a map $\nu: P \rightarrow \mathbf{N}$, we set $T^\nu := \prod_{x \in P} T_x^{\nu(x)}$.

(2.3) Let H be a finite distributive lattice. A join-irreducible element in H is an element $\alpha \in H$ such that α can not be expressed as a join of two elements different from α . That is, $\alpha = \beta \vee \gamma \Rightarrow \alpha = \beta$ or $\alpha = \gamma$. Note that we treat the unique minimal element of H as a join-irreducible element.

Let P be the set of all join-irreducible elements in H . Then it is known that H is isomorphic to $J(P) \setminus \{\emptyset\}$ ordered by inclusion, where $J(P)$ is the set of all poset ideals of P . The isomorphisms $\Phi: H \rightarrow J(P) \setminus \{\emptyset\}$ and $\Psi: J(P) \setminus \{\emptyset\} \rightarrow H$ are given by

$$\begin{aligned} \Phi(\alpha) &:= \{x \in P \mid x \leq \alpha \text{ in } H\} \quad \text{for } \alpha \in H \text{ and} \\ \Psi(I) &:= \bigvee_{x \in I} x \quad \text{for } I \in J(P) \setminus \{\emptyset\}. \end{aligned}$$

(2.4) Let L be another distributive lattice, Q the set of join-irreducible elements in L and $\varphi: H \rightarrow L$ a surjective lattice homomorphism. We set

$$\varphi^*(\beta) := \bigwedge_{\alpha \in \varphi^{-1}(\beta)} \alpha.$$

Then

Lemma 2.5 (1) $\varphi^*(\varphi(\alpha)) \leq \alpha$ for any $\alpha \in H$.

(2) $\varphi(\varphi^*(\beta)) = \beta$ for any $\beta \in L$.

(3) $\beta_1 \leq \beta_2 \Rightarrow \varphi^*(\beta_1) \leq \varphi^*(\beta_2)$.

(4) For $\alpha \in H$ and $\beta \in L$, $\varphi(\alpha) \geq \beta \iff \alpha \geq \varphi^*(\beta)$.

(5) $\varphi^*(\beta_1 \vee \beta_2) = \varphi^*(\beta_1) \vee \varphi^*(\beta_2)$.

(6) $y \in Q \Rightarrow \varphi^*(y) \in P$.

(7) If y_0 is the unique minimal element of L , then $\varphi^*(y_0)$ is the unique minimal element of H .

Proof (1): Since $\alpha \in \varphi^{-1}(\varphi(\alpha))$, we see that

$$\alpha \geq \bigwedge_{\alpha' \in \varphi^{-1}(\varphi(\alpha))} \alpha' = \varphi^*(\varphi(\alpha)).$$

(2): Since φ is a lattice homomorphism, it holds

$$\varphi(\varphi^*(\beta)) = \varphi\left(\bigwedge_{\alpha \in \varphi^{-1}(\beta)} \alpha\right) = \bigwedge_{\alpha \in \varphi^{-1}(\beta)} \varphi(\alpha) = \beta.$$

(3): Take $\alpha_0 \in \varphi^{-1}(\beta_1)$. Then for any $\alpha \in \varphi^{-1}(\beta_2)$, $\varphi(\alpha_0 \wedge \alpha) = \varphi(\alpha_0) \wedge \varphi(\alpha) = \beta_1 \wedge \beta_2 = \beta_1$. Therefore, $\alpha_0 \wedge \alpha \in \varphi^{-1}(\beta_1)$ so $\varphi^*(\beta_1) = \bigwedge_{\alpha' \in \varphi^{-1}(\beta_1)} \alpha' \leq \alpha_0 \wedge \alpha \leq \alpha$. Since α is an arbitrary element of $\varphi^{-1}(\beta_2)$, we see that

$$\varphi^*(\beta_1) \leq \bigwedge_{\alpha \in \varphi^{-1}(\beta_2)} \alpha = \varphi^*(\beta_2).$$

(4): If $\varphi(\alpha) \geq \beta$, then $\alpha \geq \varphi^*(\varphi(\alpha)) \geq \varphi^*(\beta)$ by (1) and (3). On the other hand, if $\alpha \geq \varphi^*(\beta)$, then $\varphi(\alpha) \geq \varphi(\varphi^*(\beta)) = \beta$ by (2).

(5): First we see, by (3), that

$$\varphi^*(\beta \vee \gamma) \geq \varphi^*(\beta) \vee \varphi^*(\gamma).$$


On the other hand, $\varphi(\varphi^*(\beta) \vee \varphi^*(\gamma)) = \varphi(\varphi^*(\beta)) \vee \varphi(\varphi^*(\gamma)) = \beta \vee \gamma$ by (2). Therefore $\varphi^*(\beta) \vee \varphi^*(\gamma) \in \varphi^{-1}(\beta \vee \gamma)$, and we see that

$$\varphi^*(\beta \vee \gamma) = \bigwedge_{\alpha \in \varphi^{-1}(\beta \vee \gamma)} \alpha \leq \varphi^*(\beta) \vee \varphi^*(\gamma).$$

(6): If $\varphi^*(y) = \alpha \vee \alpha'$, then $\varphi(\alpha) \vee \varphi(\alpha') = \varphi(\alpha \vee \alpha') = \varphi(\varphi^*(y)) = y$. So $y = \varphi(\alpha)$ or $y = \varphi(\alpha')$, since y is join-irreducible. By symmetry, we may assume that $y = \varphi(\alpha)$. Then $\varphi^*(y) = \varphi^*(\varphi(\alpha)) \leq \alpha$ by (1). On the other hand, $\varphi^*(y) = \alpha \vee \alpha' \geq \alpha$. Therefore $\varphi^*(y) = \alpha$.

(7): Clear. ■

Remark 2.6 $\varphi^*(\beta_1) \wedge \varphi^*(\beta_2) \neq \varphi^*(\beta_1) \wedge \varphi^*(\beta_2)$ in general. For example, let



and set $\varphi(\alpha_i) = \beta_i$ for $i = 1, 2, 3, 4$ and $\varphi(-\infty) = \beta_1$. Then $\varphi^*(\beta_i) = \alpha_i$ for $i = 2, 3$ and $\varphi^*(\beta_1) = -\infty$. Therefore, $\varphi^*(\beta_2 \wedge \beta_3) = \varphi^*(\beta_1) = -\infty \neq \alpha_1 = \varphi^*(\beta_2) \wedge \varphi^*(\beta_3)$.

By (2) and (3) of Lemma 2.5, we may regard L as a subposet (not a sublattice) of H by φ^* . Then, by (6) of Lemma 2.5, Q is identified as a subset of P . Note that Q contains the unique minimal element of H under this identification by (7) of Lemma 2.5.

Lemma 2.7 *By the identification above, the composition map $J(P) \setminus \{\emptyset\} \simeq H \xrightarrow{\varphi} L \simeq J(Q) \setminus \{\emptyset\}$ is identical to the map $I \mapsto I \cap Q$.*

Proof Recall that lattice isomorphisms $H \simeq J(P) \setminus \{\emptyset\}$ and $L \simeq J(Q) \setminus \{\emptyset\}$ are obtained by $\Phi_H: H \rightarrow J(P) \setminus \{\emptyset\}$, $(\alpha \mapsto \{x \in P \mid x \leq \alpha\})$ and $\Phi_L: L \rightarrow J(Q) \setminus \{\emptyset\}$, $(\beta \mapsto \{y \in Q \mid y \leq \beta\})$. Let α be an arbitrary element of H . Then $\Phi_L(\varphi(\alpha)) = \{y \in Q \mid y \leq \varphi(\alpha)\}$. Since $y \leq \varphi(\alpha)$ if and only if $\varphi^*(y) \leq \alpha$ by (4) of Lemma 2.5, we see that by the embedding of L in H by φ^* , $\Phi_L(\varphi(\alpha))$ is identified to $\{y \in Q \mid y \leq \alpha\}$ which is equal to $\{x \in P \mid x \leq \alpha\} \cap Q = \Phi_H(\alpha) \cap Q$. ■

2.3 Determinantal rings

(2.5) Let m, n be integers with $0 < m \leq n$ and K a field. For an $m \times n$ matrix M with entries in a K algebra S , we denote by M^T the transposed matrix of M , by $I_t(M)$ the ideal of S generated by the t -minors of M , by $M^{\leq i}$ the $i \times n$ matrix consisting of first i -rows of M , by $M_{\leq j}$ the $m \times j$ matrix consisting of first j -columns of M , by $\Gamma(M)$ the set of maximal minors of M and by $K[M]$ the K -subalgebra of S generated by the entries of M .

We define posets $\Gamma(m \times n)$ and $\Gamma'(m \times n)$ by $\Gamma(m \times n) := \{[c_1, \dots, c_m] \mid 1 \leq c_1 < \dots < c_m \leq n\}$ and $\Gamma'(m \times n) := \{[c_1, \dots, c_r] \mid r \leq m, 1 \leq c_1 < \dots < c_r \leq n\}$. The order of $\Gamma'(m \times n)$ is defined by

$$\begin{aligned} [c_1, \dots, c_r] &\leq [d_1, \dots, d_s] \\ \xLeftrightarrow{\text{def}} \quad &r \geq s, \quad c_i \leq d_i \text{ for } i = 1, 2, \dots, s, \end{aligned}$$

and the order of $\Gamma(m \times n)$ is defined by that of $\Gamma'(m \times n)$. Note that $\Gamma'(m \times n)$ is a distributive lattice and $\Gamma(m \times n)$ is a sublattice of $\Gamma'(m \times n)$. For $[c_1, \dots, c_r] \in \Gamma'(m \times n)$, we define its size to be r and denote it by $\text{size}[c_1, \dots, c_r]$.

We also define the poset $\Delta(m \times n)$ by $\Delta(m \times n) := \{[\alpha|\beta] \mid \alpha, \beta \in \Gamma'(m \times n), \text{size}\alpha = \text{size}\beta\}$ and define the order of $\Delta(m \times n)$ by

$$\begin{aligned} & [\alpha|\beta] \leq [\alpha'|\beta'] \\ \stackrel{\text{def}}{\iff} & \alpha \leq \alpha' \text{ in } \Gamma'(m \times m) \text{ and } \beta \leq \beta' \text{ in } \Gamma'(m \times n). \end{aligned}$$

Note also that $\Delta(m \times n)$ is a distributive lattice.

For $\gamma \in \Gamma(m \times n)$, we set $\Gamma(m \times n; \gamma) := \{\delta \in \Gamma(m \times n) \mid \delta \geq \gamma\}$. $\Gamma'(m \times n; \gamma)$ and $\Delta(m \times n; \delta)$ are defined similarly.

For an $m \times n$ matrix $M = (m_{ij})$ and $\delta = [\alpha|\beta] = [c_1, \dots, c_r | d_1, \dots, d_r] \in \Delta(m \times n)$, we denote the minor $\det(m_{c_i d_j})$ by δ_M or $[\alpha|\beta]_M$. We also denote the maximal minor $\det(m_{i c_j})$ of M by δ_M or $[c_1, \dots, c_m]_M$, where $\delta = [c_1, \dots, c_m] \in \Gamma(m \times n)$.

We use standard terminology on Hodge algebras. Standard references on the notion of Hodge algebra are [DEP2] and [BH, Chapter 7]. However, we use the term “algebra with straightening law” (ASL for short) to mean an ordinal Hodge algebra in the terminology of [DEP2].

Let $X = (X_{ij})$ be an $m \times n$ matrix of indeterminates, i.e., entries X_{ij} are independent indeterminates. Then the following facts are known.

Theorem 2.8 ([DEP1]) (1) $K[\Gamma(X)]$ is an ASL over K generated by $\Gamma(m \times n)$ with structure map $\delta \mapsto \delta_X$.

(2) $K[X]$ is an ASL over K generated by $\Delta(m \times n)$ with structure map $[\alpha|\beta] \mapsto [\alpha|\beta]_X$.

(2.6) Now let V be an n -dimensional K -vector space. Then the set of m -dimensional subspaces of V has a structure of an algebraic variety, called the Grassmann variety. We denote this variety by $G_m(V)$. It is known that if $X = (X_{ij})$ is the $m \times n$ matrix of indeterminates, then $K[\Gamma(X)]$ is the homogeneous coordinate ring of $G_m(V)$.

Now fix a complete flag $0 = V_0 \subsetneq V_1 \subsetneq \dots \subsetneq V_n = V$ of subspaces of V .

Definition 2.9 For $[a_1, \dots, a_m] \in \Gamma(m \times n)$, we define $\Omega(a_1, \dots, a_m) := \{W \in G_m(V) \mid \dim(W \cap V_{a_i}) \geq i \text{ for } i = 1, 2, \dots, m\}$.

It is known that $\Omega(a_1, \dots, a_m)$ is subvariety of $G_m(V)$ and called the Schubert subvariety of $G_m(V)$.

Set $b_i := n - a_{m-i+1} + 1$ for $i = 1, \dots, m$ and $\gamma := [b_1, b_2, \dots, b_m] \in \Gamma(m \times n)$. It is known that there is the universal $m \times n$ matrix Z_γ with $I_i((Z_\gamma)_{\leq b_i-1}) = (0)$ for $i = 1, \dots, m$. That is,

(1) $I_i((Z_\gamma)_{\leq b_i-1}) = (0)$ for $i = 1, \dots, m$ and

- (2) if $I_i(M_{\leq b_i-1}) = (0)$ for $i = 1, \dots, m$ for a matrix with entries in some k -algebra S , then there is a unique k -algebra homomorphism $k[Z_\gamma] \rightarrow S$ which maps Z_γ to M .

And it is also known that $K[\Gamma(Z_\gamma)]$ is the homogeneous coordinate ring of $\Omega(a_1, \dots, a_m)$.

From now on, until the end of this paper, we fix $\gamma = [b_1, b_2, \dots, b_m] \in \Gamma(m \times n)$. Set $\gamma^* = [1, \dots, m | b_1, \dots, b_m]$ and $R = K[X]/(\Delta(m \times n) \setminus \Delta(m \times n; \gamma^*))K[X]$. Then it is easy to see that the image of X in R satisfies the universal property above.

(2.7) Set

$$W := \begin{pmatrix} W_{11} & W_{12} & \cdots & W_{1m} \\ W_{21} & W_{22} & \cdots & W_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ W_{m1} & W_{m2} & \cdots & W_{mm} \end{pmatrix}$$

and

$$U_\gamma := \begin{pmatrix} 0 & \cdots & 0 & U_{1b_1} & \cdots & U_{1b_2-1} & U_{1b_2} & \cdots & \cdots & U_{1b_m} & \cdots & U_{1n} \\ 0 & \cdots & 0 & 0 & \cdots & 0 & U_{2b_2} & \cdots & \cdots & U_{2b_m} & \cdots & U_{2n} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & U_{mb_m} & \cdots & U_{mn} \end{pmatrix},$$

where W_{ij} and U_{ij} are independent indeterminates. Then

Theorem 2.10 ([Miy]) *The natural map $K[X]/(\Delta(m \times n) \setminus \Delta(m \times n; \gamma^*))K[X] \rightarrow K[WU_\gamma]$ induced by $K[X] \rightarrow K[WU_\gamma]$, ($X \mapsto WU_\gamma$) is an isomorphism. In particular, WU_γ has the universal property above.*

So we set $Z_\gamma := WU_\gamma$ in the following.

Here we state a result on the initial algebra of $K[WU_\gamma]$ in $K[W, U_\gamma]$ along the same line as [BRW, section 3] (see also [BC]). As a by-product, we obtain another proof of Theorem 2.10.

First we introduce a diagonal monomial order (i.e. a monomial order such that leading monomial of any minor of W or U_γ is the product of the entries of its main diagonal) on $K[W, U_\gamma]$ with $W_{ij} > W_{i'j}$, $W_{ij} > W_{ij'}$, $U_{ij} > U_{i'j}$ and $U_{ij} > U_{ij'}$ for any possible i, i', j and j' with $i < i'$ and $j < j'$. For example degree lexicographic order on the polynomial ring $K[W, U_\gamma]$ defined by $W_{11} > W_{21} > \cdots > W_{m1} > W_{12} > \cdots > W_{mm} > U_{1b_1} > U_{1b_1+1} > \cdots > U_{1n} > U_{2b_2} > \cdots > U_{mn}$. Then we obtain the following

Lemma 2.11 (1) For $[c_1, \dots, c_r | d_1, \dots, d_r] \in \Delta(m \times n; \gamma^*)$,

$$\text{lm}([c_1, \dots, c_r | d_1, \dots, d_r]_{WU_\gamma}) = \prod_{j=1}^r W_{c_j j} U_{j d_j}.$$

(2) If $\mu = \prod_{i=1}^u [c_{i1}, \dots, c_{ir(i)} | d_{i1}, \dots, d_{ir(i)}]$ is a standard monomial on $\Delta(m \times n; \gamma^*)$, then

$$\text{lm}(\mu_{WU_\gamma}) = \prod_{i=1}^u \prod_{j=1}^{r(i)} W_{c_{ij}j} U_{jd_{ij}},$$

where $\mu_{WU_\gamma} := \prod_{i=1}^u [c_{i1}, \dots, c_{ir(i)} | d_{i1}, \dots, d_{ir(i)}]_{WU_\gamma}$.

(3) If μ and μ' are standard monomials on $\Delta(m \times n; \gamma^*)$ with $\mu \neq \mu'$, then $\text{lm}(\mu_{WU_\gamma}) \neq \text{lm}(\mu'_{WU_\gamma})$. In particular, $\{\mu_{WU_\gamma} \mid \mu \text{ is a standard monomial on } \Delta(m \times n; \gamma^*)\}$ is linearly independent over K .

(4) The natural map $K[X]/(\Delta(m \times n) \setminus \Delta(m \times n; \gamma^*))K[X] \rightarrow K[WU_\gamma]$ induced by $K[X] \rightarrow K[WU_\gamma]$, $(X \mapsto WU_\gamma)$ is an isomorphism.

(5) For any $a \in K[WU_\gamma]$, there is unique standard monomial μ on $\Delta(m \times n; \gamma^*)$ such that $\text{lm}(a) = \text{lm}(\mu_{WU_\gamma})$.

Proof (1) is proved by the same way as the proof of [BRW, Lemma 3.1] or by using the following equation on determinants.

$$\begin{aligned} & [c_1, \dots, c_r | d_1, \dots, d_r]_{WU_\gamma} \\ &= \sum_{1 \leq j_1 < \dots < j_r \leq m} [c_1, \dots, c_r | j_1, \dots, j_r]_W [j_1, \dots, j_r | d_1, \dots, d_r]_{U_\gamma}. \end{aligned}$$

The rest is proved by the same way as [BRW, Section 3] or by using the ASL structure of $K[X]/(\Delta(m \times n) \setminus \Delta(m \times n; \gamma^*))K[X]$. ■

As a corollary, we see the following

Proposition 2.12 $Z_\gamma = WU_\gamma$ has the universal property and $\{[\alpha|\beta]_{Z_\gamma} \mid [\alpha|\beta] \in \Delta(m \times n; \gamma^*)\}$ is a sagbi basis of $K[Z_\gamma]$.

(2.8) We set $D_{m,n} := \{(\alpha, \beta) \mid \alpha, \beta \in \Gamma'(m \times n), \text{size}\alpha = \text{size}\beta\}$ and $D'_{m,n} := D_{m-1,n} \cup \Gamma(m \times n)$. We define the order on $D_{m,n}$ by lexicographic way, that is,

$$(\alpha, \beta) < (\alpha', \beta') \stackrel{\text{def}}{\iff} \alpha < \alpha' \text{ or } \alpha = \alpha', \beta < \beta'.$$

We also define the order on $D'_{m,n}$ by extending the orders of $D_{m-1,n}$ and $\Gamma(m \times n)$ and set

$$\delta < (\alpha, \beta) \stackrel{\text{def}}{\iff} \delta < \alpha \text{ in } \Gamma'(m \times n)$$

for $\delta \in \Gamma(m \times n)$ and $(\alpha, \beta) \in D_{m-1,n}$. Then

Theorem 2.13 ([DP, Section 5]) (1) $K[X^T X]$ is a Hodge algebra over K generated by $D_{m,n}$ with structure map $(\alpha, \beta) \mapsto [\alpha|\beta]_{X^T X}$, where the ideal of monomials on $D_{m,n}$ defining this Hodge algebra structure is generated by

$$\{(\alpha, \beta)(\alpha', \beta') \mid \beta \not\leq \alpha' \text{ and } \beta' \not\leq \alpha\}.$$

(2) $K[X^T X, \Gamma(X)]$ is a Hodge algebra over K generated by $D'_{m,n}$ with structure map $\delta \mapsto \delta_X$ for $\delta \in \Gamma(m \times n)$ and $(\alpha, \beta) \mapsto [\alpha|\beta]_{X^T X}$ for $(\alpha, \beta) \in D_{m-1,n}$, where the ideal of monomials on $D'_{m,n}$ defining this Hodge algebra structure is generated by

$$\begin{aligned} & \{(\alpha, \beta)(\alpha', \beta') \mid (\alpha, \beta), (\alpha', \beta') \in D_{m-1,n}, \beta \not\leq \alpha' \text{ and } \beta' \not\leq \alpha\} \\ \cup & \ \{\delta(\alpha, \beta) \mid \delta \in \Gamma(m \times n), (\alpha, \beta) \in D_{m-1,n} \text{ and } \delta \not\leq \alpha \text{ in } \Gamma'(m \times n)\} \\ \cup & \ \{\delta\delta' \mid \delta, \delta' \in \Gamma(m \times n), \delta \not\sim \delta'\}. \end{aligned}$$

In fact, DeConcini-Procesi [DP, Lemmas 5.2, 5.3 and 5.4] essentially proved the following fact also.

Theorem 2.14 Let $T = (T_{ij})$ be the $n \times n$ symmetric matrix of indeterminates, i.e., $\{T_{ij}\}_{1 \leq i \leq j \leq n}$ are independent indeterminates and $T_{ji} = T_{ij}$ for $j > i$. Then $K[T]$ is a Hodge algebra over K generated by $D_{n,n}$ with structure map $(\alpha, \beta) \mapsto [\alpha|\beta]_T$.

For a standard monomial $\nu = \prod_{i=1}^v ([c_{i1}, \dots, c_{ir(i)}], [d_{i1}, \dots, d_{ir(i)}])$ on $D_{m,n}$, we set $\nu_X := \prod_{i=1}^v [c_{i1}, \dots, c_{ir(i)} | d_{i1}, \dots, d_{ir(i)}]_{X^T X}$ and for a standard monomial $\nu' = \prod_{i=1}^{v'} [e_{i1}, \dots, e_{im}] \prod_{i=1}^{v''} ([c_{i1}, \dots, c_{ir(i)}], [d_{i1}, \dots, d_{ir(i)}])$ on $D'_{m,n}$, we set $\nu'_X := \prod_{i=1}^{v'} [e_{i1}, \dots, e_{im}]_X \prod_{i=1}^{v''} [c_{i1}, \dots, c_{ir(i)} | d_{i1}, \dots, d_{ir(i)}]_{X^T X}$.

By the same argument as the proof of Lemma 2.11, we see the following results.

Lemma 2.15 (1) For $[c_1, \dots, c_r | d_1, \dots, d_r] \in \Delta(m \times m)$, we have

$$\text{lm}([c_1, \dots, c_r | d_1, \dots, d_r]_{W^T W}) = \prod_{j=1}^r W_{jc_j} W_{jd_j}.$$

(2) If $\nu = \prod_{i=1}^v ([c_{i1}, \dots, c_{ir(i)}], [d_{i1}, \dots, d_{ir(i)}])$ is a standard monomial on $D_{m,m}$, then

$$\text{lm}(\nu_W) = \prod_{i=1}^v \prod_{j=1}^{r(i)} W_{jc_{ij}} W_{jd_{ij}}.$$

- (3) If ν and ν' are standard monomials on $D_{m,m}$ with $\nu \neq \nu'$, then $\text{lm}(\nu_W) \neq \text{lm}(\nu'_W)$. In particular, $\{\nu_W \mid \nu \text{ is a standard monomial on } D_{m,m}\}$ is linearly independent over K .
- (4) Let $T' = (T_{ij})$ be the $m \times m$ symmetric matrix of indeterminates. Then the K -algebra homomorphism $K[T'] \rightarrow K[W^T W]$, $(T' \mapsto W^T W)$ is an isomorphism.
- (5) For any $a \in K[W^T W]$, there is a unique standard monomial ν on $D_{m,m}$ such that $\text{lm}(a) = \text{lm}(\nu_W)$. In particular, $\{[\alpha|\beta]_{W^T W} \mid (\alpha, \beta) \in D_{m,m}\}$ is a sagbi basis of $K[W^T W]$.

Lemma 2.16 (1) If $\nu = [1, \dots, m]^{v'} \prod_{i=1}^{v''} ([c_{i1}, \dots, c_{ir(i)}], [d_{i1}, \dots, d_{ir(i)}])$ is a standard monomial on $D'_{m,m}$, then

$$\text{lm}(\nu_W) = \left(\prod_{j=1}^m W_{jj} \right)^{v'} \prod_{i=1}^{v''} \prod_{j=1}^{r(i)} W_{jc_{ij}} W_{jd_{ij}}.$$

- (2) If ν and ν' are standard monomials on $D'_{m,m}$ with $\nu \neq \nu'$, then $\text{lm}(\nu_W) \neq \text{lm}(\nu'_W)$. In particular, $\{\nu_W \mid \nu \text{ is a standard monomial on } D'_{m,m}\}$ is linearly independent over K .
- (3) For any $a \in K[W^T W, \det W]$, there is a unique standard monomial ν on $D'_{m,m}$ such that $\text{lm}(a) = \text{lm}(\nu_W)$. In particular, $\{\det W\} \cup \{[\alpha|\beta]_{W^T W} \mid (\alpha, \beta) \in D_{m-1,m}\}$ is a sagbi basis of $K[W^T W, \det W]$.

Finally in this section we make the following remark. Let T be the $n \times n$ symmetric matrix of indeterminates. Set

$$R = K[T] / \{(\alpha, \beta) \in D_{n,n} \mid \alpha \not\geq \gamma\} K[T],$$

where $\gamma = [b_1, \dots, b_m] \in \Gamma(m \times n)$ is the fixed element. Then R is a Hodge algebra over K generated by $D_\gamma := \{(\alpha, \beta) \in D_{n,n} \mid \alpha \geq \gamma\}$, since $D_{n,n} \setminus D_\gamma$ is a poset ideal of $D_{n,n}$. Let T_γ be the image of T in R . Then T_γ has the following universal property, where we set $b_{m+1} := n + 1$.

- (1) $I_i((T_\gamma)_{\leq b_i-1}) = (0)$ for $i = 1, \dots, m + 1$ and
- (2) if M is an $n \times n$ symmetric matrix with entries in some K -algebra S such that $I_i(M_{\leq b_i-1}) = (0)$ for $i = 1, \dots, m + 1$, then there is a unique K -algebra homomorphism $K[T_\gamma] \longrightarrow S$ which maps T_γ to M .

By the same argument as the proof of Lemma 2.11, we see the following result, cf. [BRW, Remark 3.10].

Lemma 2.17 (1) For $([c_1, \dots, c_r], [d_1, \dots, d_r]) \in D_\gamma$,

$$\text{lm}([c_1, \dots, c_r | d_1, \dots, d_r]_{Z_\gamma^T Z_\gamma}) = \prod_{j=1}^r W_{jj}^2 U_{jc_j} U_{jd_j}.$$

(2) If $\nu = \prod_{i=1}^l ([c_{i1}, \dots, c_{ir(i)}], [d_{i1}, \dots, d_{ir(i)}])$ is a standard monomial on D_γ , then

$$\text{lm}(\nu_{Z_\gamma}) = \prod_{i=1}^l \prod_{j=1}^{r(i)} W_{jj}^2 U_{jc_{ij}} U_{jd_{ij}}.$$

(3) If ν and ν' are standard monomials on D_γ with $\nu \neq \nu'$, then $\text{lm}(\nu_{Z_\gamma}) \neq \text{lm}(\nu'_{Z_\gamma})$. In particular, $\{\nu_{Z_\gamma} \mid \nu \text{ is a standard monomial on } D_\gamma\}$ is linearly independent over K .

(4) The natural map $K[T]/(D_{n,n} \setminus D_\gamma)K[T] \rightarrow K[Z_\gamma^T Z_\gamma]$ induced by $K[T] \rightarrow K[Z_\gamma^T Z_\gamma]$, $(T \mapsto Z_\gamma^T Z_\gamma)$ is an isomorphism.

(5) $Z_\gamma^T Z_\gamma$ is the universal $n \times n$ symmetric matrix with $I_i((M)_{\leq b_i-1}) = (0)$ for $i = 1, \dots, m+1$ and $\{[\alpha|\beta]_{Z_\gamma^T Z_\gamma} \mid (\alpha, \beta) \in D_\gamma\}$ is a sagbi basis of $K[Z_\gamma^T Z_\gamma]$.

3 Rings of invariants and sagbi bases

Recall that we have fixed integers m, n with $0 < m \leq n$ and $\gamma = [b_1, b_2, \dots, b_m] \in \Gamma(m \times n)$.

For any $g \in \text{GL}(m, K)$, $I_i((gZ_\gamma)_{\leq b_i-1}) = (0)$ for $i = 1, \dots, m$. So, by the universal property of Z_γ , there is a K -automorphism of $K[Z_\gamma]$ sending Z_γ to gZ_γ , i.e., any subgroup of $\text{GL}(m, K)$ acts on $K[Z_\gamma]$.

Definition 3.1 ([DP, Section 2]) $f \in K[Z_\gamma]$ is called an absolute $\text{O}(m)$ -invariant if f is an $\text{O}(m, B)$ -invariant in $B[Z_\gamma]$ for any K -algebra B . The set of absolute $\text{O}(m)$ invariants is denoted by $K[Z_\gamma]^{\text{O}(m, -)}$. Absolute $\text{SO}(m)$ invariants and $K[Z_\gamma]^{\text{SO}(m, -)}$ are defined similarly.

First we recall the following results.

Theorem 3.2 ([DP, Section 5]) Let X an $m \times n$ matrix of indeterminates. Then

$$(1) K[X]^{\text{O}(m,-)} = K[X^T X].$$

$$(2) K[X]^{\text{SO}(m,-)} = K[X^T X, \Gamma(X)].$$

By considering the case where $m = n$, we see the following

$$\textbf{Corollary 3.3} \quad (1) K[W]^{\text{O}(m,-)} = K[W^T W].$$

$$(2) K[W]^{\text{SO}(m,-)} = K[W^T W, \det W].$$

Here we note the following basic fact.

Lemma 3.4 *Let $K[X_1, \dots, X_m, Y_1, \dots, Y_n]$ be a polynomial ring over K with monomial order and A a K -subalgebra of $K[X_1, \dots, X_m]$. If S is a sagbi basis of A in $K[X_1, \dots, X_m]$, then $S \cup \{Y_1, \dots, Y_n\}$ is a sagbi basis of $A[Y_1, \dots, Y_n]$ in $K[X_1, \dots, X_m, Y_1, \dots, Y_n]$.*

Now we state the following

Theorem 3.5 (1) $\{[\alpha|\beta]_{Z_\gamma^T Z_\gamma} \mid \alpha, \beta \in \Gamma'(m \times n; \gamma), \alpha \leq \beta, \text{size}\alpha = \text{size}\beta\}$ is a sagbi basis of $K[W^T W, U_\gamma] \cap K[Z_\gamma]$. In particular, $K[Z_\gamma]^{\text{O}(m,-)} = K[W^T W, U_\gamma] \cap K[Z_\gamma] = K[Z_\gamma^T Z_\gamma]$.

(2) $\{[\alpha|\beta]_{Z_\gamma^T Z_\gamma} \mid \alpha, \beta \in \Gamma'(m \times n; \gamma), \alpha \leq \beta, \text{size}\alpha = \text{size}\beta < m\} \cup \{\delta_{Z_\gamma} \mid \delta \in \Gamma(m \times n; \gamma)\}$ is a sagbi basis of $K[W^T W, \det W, U_\gamma] \cap K[Z_\gamma]$. In particular, $K[Z_\gamma]^{\text{SO}(m,-)} = K[W^T W, \det W, U_\gamma] \cap K[Z_\gamma] = K[Z_\gamma^T Z_\gamma, \Gamma(Z_\gamma)]$.

Proof (1) It is clear that $K[W^T W, U_\gamma] \cap K[Z_\gamma] \supset K[Z_\gamma^T Z_\gamma]$. So $[\alpha|\beta]_{Z_\gamma^T Z_\gamma} \in K[W^T W, U_\gamma] \cap K[Z_\gamma]$ for any $\alpha, \beta \in \Gamma'(m \times n; \gamma)$ with $\text{size}\alpha = \text{size}\beta$. Let a be an arbitrary element of $K[W^T W, U_\gamma] \cap K[Z_\gamma]$. Then by Lemma 2.11, we see that there is unique standard monomial μ on $\Delta(m \times n; \gamma^*)$ such that $\text{lm}(a) = \text{lm}(\mu_{Z_\gamma})$.

Set $\mu = \prod_{i=1}^u [\alpha_i|\beta_i]$, $\alpha_1 \leq \dots \leq \alpha_u$, $\beta_1 \leq \dots \leq \beta_u$, and $\alpha_i = [c_{i1}, \dots, c_{ir(i)}]$, $\beta_i = [d_{1i}, \dots, d_{ir(i)}]$ for $i = 1, \dots, u$. Then

$$\text{lm}(a) = \text{lm}(\mu_{Z_\gamma}) = \prod_{i=1}^u \prod_{j=1}^{r(i)} W_{c_{ij}j} U_{jd_{ij}}. \quad (3.1)$$

On the other hand, we see, by Lemmas 2.15 and 3.4, that

$$\text{lm}(a) = \left(\prod_{i=1}^v \prod_{j=1}^{s(i)} W_{jc'_{ij}} W_{jd'_{ij}} \right) \times (\text{a monomial of } U'_{\cdot} s) \quad (3.2)$$

for some c'_{ij} and d'_{ij} , since $a \in K[W^T W, U_\gamma]$.

By comparing (3.1) and (3.2), we see that $c_{ij} = j$, $c'_{ij} = j$, and $d'_{ij} = j$ for any possible i, j , u is even and $r(2i-1) = r(2i)$ for $i = 1, \dots, u/2$. So if we set $\alpha'_i = [d_{2i-1,1}, \dots, d_{2i-1,r(2i)}]$ and $\beta'_i = [d_{2i,1}, \dots, d_{2i,r(2i)}]$, then $\alpha'_i, \beta'_i \in \Gamma'(m \times n; \gamma)$ for $i = 1, \dots, u/2$, $\prod_{i=1}^{u/2} (\alpha'_i, \beta'_i)$ is a standard monomial on $D_{m,n}$ and

$$\text{lm}(a) = \prod_{i=1}^{u/2} \text{lm}([\alpha'_i | \beta'_i]_{Z_\gamma^T Z_\gamma}).$$

The former part of (1) follows. Since the action of $O(m)$ on $K[Z_\gamma]$ is induced by the action of $O(m)$ on $K[W]$, we see the latter part of (1).

(2) Let a be an arbitrary element of $K[W^T W, \det W, U_\gamma] \cap K[Z_\gamma]$. By the same argument as in the proof of (1), we see that there is a standard monomial μ on $\Delta(m \times n; \gamma^*)$ such that $\text{lm}(a) = \text{lm}(\mu_{Z_\gamma})$. Set, as before, $\mu = \prod_{i=1}^u [\alpha_i | \beta_i]$, $\alpha_1 \leq \dots \leq \alpha_u$, $\beta_1 \leq \dots \leq \beta_u$, and $\alpha_i = [c_{i1}, \dots, c_{ir(i)}]$, $\beta_i = [d_{i1}, \dots, d_{ir(i)}]$ for $i = 1, \dots, u$. Then by using Lemma 2.16 and the same argument as in the proof of (1), we see that $c_{ij} = j$ for any i, j and there is $w \in \mathbf{N}$ such that $w \leq u$, $r(i) = m \iff i \leq w$, $u-w$ is even and $r(2i-1+w) = r(2i+w)$ for $i = 1, \dots, (u-w)/2$. So if we set $\alpha''_i = [d_{2i-1+w,1}, \dots, d_{2i-1+w,r(2i+w)}]$ and $\beta''_i = [d_{2i+w,1}, \dots, d_{2i+w,r(2i+w)}]$ for $i = 1, \dots, (u-w)/2$ and $\delta_i = [d_{i1}, \dots, d_{im}]$ for $i = 1, \dots, w$, then

$$\text{lm}(a) = \prod_{i=1}^u \prod_{j=1}^{r(i)} W_{jj} U_{jd_{ij}} = \prod_{i=1}^w \text{lm}((\delta_i)_{Z_\gamma}) \prod_{i=1}^{(u-w)/2} \text{lm}([\alpha''_i | \beta''_i]_{Z_\gamma^T Z_\gamma}).$$

The rest is proved as (1). ■

4 Doset Hibi rings and generalization

Let H be a finite distributive lattice, P the set of join-irreducible elements, $\{T_x\}_{x \in P}$ a family of indeterminates indexed by P . Recall that $\overline{\mathcal{T}}(P)$ is the set of order reversing maps from P to $\mathbf{N}$ and $\mathcal{T}(P)$ is the set of strictly order reversing maps from P to $\mathbf{N} \setminus \{0\}$. Recall also $T^\nu = \prod_{x \in P} T_x^{\nu(x)}$ for $\nu \in \overline{\mathcal{T}}(P)$.

Hibi [Hib] defined the ring $\mathcal{R}_K(H)$, called the Hibi ring nowadays, as follows.

Definition 4.1 $\mathcal{R}_K(H) := K[\prod_{x \in \alpha} T_x \mid \alpha \in H]$.

Hibi [Hib] showed the following theorems.

Theorem 4.2 $\mathcal{R}_K(H)$ is an ASL over K generated by H with structure map $\alpha \mapsto \prod_{x \leq \alpha} T_x$. The straightening law of $\mathcal{R}_K(H)$ is $\alpha\beta = (\alpha \wedge \beta)(\alpha \vee \beta)$ for $\alpha, \beta \in \bar{H}$ with $\alpha \not\leq \beta$.

Theorem 4.3 $\mathcal{R}_K(H)$ is a free K -module with basis $\{T^\nu \mid \nu \in \overline{\mathcal{T}}(P)\}$. In particular, by Theorem 2.1, $\mathcal{R}_K(H)$ is a normal affine semigroupring and hence Cohen-Macaulay.

By analyzing the relations of the leading monomials of the elements of $\Delta(m \times n; \gamma^*)$, embeded by the structure map of the ASL $K[Z_\gamma]$ by Lemma 2.11, we see the following

Proposition 4.4 in $K[Z_\gamma]$ is the Hibi ring over K generated by $\Delta(m \times n; \gamma^*)$. In particular, it is normal and therefore, by [CHV, Corollary 2.3], $K[Z_\gamma]$ is normal and Cohen-Macaulay and has rational singularities if $\text{char} K = 0$ and is F -rational if $\text{char} K > 0$.

Let us recall the definition of dosets [DEP2, Section 18].

Definition 4.5 A subset D of $H \times H$ is a doset of H if

- (1) $\{(\alpha, \alpha) \mid \alpha \in H\} \subset D \subset \{(\alpha, \beta) \in H \times H \mid \alpha \leq \beta\}$ and
- (2) for α, β and $\gamma \in H$ with $\alpha \leq \beta \leq \gamma$,

$$(\alpha, \gamma) \in D \iff (\alpha, \beta) \in D, (\beta, \gamma) \in D.$$

Now let L be another distributive lattice Q the set of join-irreducible elements of L and $\varphi: H \rightarrow L$ a surjective lattice homomorphism. We set $D := \{(\alpha, \beta) \in H \times H \mid \alpha \leq \beta, \varphi(\alpha) = \varphi(\beta)\}$. Then it is easily verified that D is a doset of H .

We define the doset Hibi ring as follows.

Definition 4.6 The doset Hibi ring over K defined by φ , denoted by $\mathcal{D}_K(\varphi)$ is the K -subalgebra of $\mathcal{R}_K(H)$ generated by $\{\alpha\beta \mid (\alpha, \beta) \in D\}$.

Note that if $(\alpha, \beta), (\alpha', \beta') \in D$, then

$$\begin{aligned} & (\alpha\beta)(\alpha'\beta') \\ &= (\alpha \wedge \alpha')(\alpha \vee \alpha')(\beta \wedge \beta')(\beta \vee \beta') \\ &= (\alpha \wedge \alpha')((\alpha \vee \alpha') \wedge (\beta \wedge \beta'))((\alpha \vee \alpha') \vee (\beta \wedge \beta'))(\beta \vee \beta') \end{aligned} \tag{4.1}$$

and

$$\begin{aligned} \varphi((\alpha \vee \alpha') \wedge (\beta \wedge \beta')) &= \varphi(\alpha \wedge \alpha') \\ \varphi((\alpha \vee \alpha') \vee (\beta \wedge \beta')) &= \varphi(\beta \vee \beta'). \end{aligned}$$

Therefore $(\alpha \wedge \alpha', (\alpha \vee \alpha') \wedge (\beta \wedge \beta')), ((\alpha \vee \alpha') \vee (\beta \wedge \beta'), \beta \vee \beta') \in D$.

In particular, by applying the straightening law repeatedly, we see that $\{\alpha_1 \alpha_2 \cdots \alpha_{2r-1} \alpha_{2r} \mid \alpha_1 \leq \alpha_2 \leq \cdots \leq \alpha_{2r}, (\alpha_{2i-1}, \alpha_{2i}) \in D \text{ for } i = 1, \dots, r\}$ is a K -free basis of the doset Hibi ring.

Example 4.7 (1) If L is a set with only one element and $\varphi: H \rightarrow L$ is the unique map, then $\mathcal{D}_K(\varphi)$ is the second Veronese subring of $\mathcal{R}_K(H)$.

(2) If $L = H$ and φ is the identity map, then $\mathcal{D}_K(\varphi) = K[\alpha^2 \mid \alpha \in H]$, which is isomorphic to $\mathcal{R}_K(H)$.

Regarding $\mathcal{R}_K(H)$ as a subring of $K[T_x \mid x \in P]$, a standard monomial $\alpha_1 \alpha_2 \cdots \alpha_s$ with $\alpha_1 \leq \alpha_2 \leq \cdots \leq \alpha_s$ corresponds to T^ν , where $\nu(x) = \#\{i \mid x \leq \alpha_i\}$. On the other hand, by the identification $L \simeq J(Q) \setminus \{\emptyset\}$ and $Q \subset P$, $\varphi(\alpha)$ corresponds to $\{y \in Q \mid y \leq \alpha \text{ in } H\}$ by Lemma 2.7. So a standard monomial $\alpha_1 \alpha_2 \cdots \alpha_{2r}$ on H with $\alpha_1 \leq \alpha_2 \leq \cdots \leq \alpha_{2r}$ satisfies $(\alpha_{2i-1}, \alpha_{2i}) \in D$ for $i = 1, \dots, r$ if and only if $\nu(y) \equiv 0 \pmod{2}$ for any $y \in Q$, where ν is an element of $\overline{\mathcal{T}}(P)$ corresponding to $\alpha_1 \cdots \alpha_{2r}$. Since Q contains the unique minimal element x_0 of H and $\nu(x_0)$ is the length of the standard monomial corresponding to $\nu \in \overline{\mathcal{T}}(P)$, we see the following

Theorem 4.8 $\mathcal{D}_K(\varphi)$ is a free K -module with basis $\{T^\nu \mid \nu \in \overline{\mathcal{T}}(P), \nu(y) \equiv 0 \pmod{2} \text{ for any } y \in Q\}$. In particular, by Theorem 2.1, $\mathcal{D}_K(\varphi)$ is a normal affine semigroup ring, and therefore, is Cohen-Macaulay.

Note the description of $\mathcal{D}_K(\varphi)$ in the theorem above depends only on H and Q . Therefore we can generalize the notion of doset Hibi ring as follows.

Definition 4.9 Let H be a finite distributive lattice, P the set of join-irreducible elements of H , Q a subset of P . We define the generalized doset Hibi ring defined by H and Q , denoted by $\mathcal{D}_K(H, Q)$, as $\mathcal{D}_K(H, Q) := K[T^\nu \mid \nu \in \overline{\mathcal{T}}(P), \nu(y) \equiv 0 \pmod{2} \text{ for any } y \in Q]$.

The following theorem is a direct consequence of the definition.

Theorem 4.10 The generalized doset Hibi ring $\mathcal{D}_K(H, Q)$ is a free K -module with basis $\{T^\nu \mid \nu \in \overline{\mathcal{T}}(P), \nu(y) \equiv 0 \pmod{2} \text{ for any } y \in Q\}$. In particular, $\mathcal{D}_K(H, Q)$ is a normal affine semigroup ring and hence is Cohen-Macaulay.

Note that if Q contains the minimal element x_0 of H , then by setting $L = J(Q) \setminus \{\emptyset\}$ and φ the map corresponding to $J(P) \setminus \{\emptyset\} \rightarrow J(Q) \setminus \{\emptyset\}$, $(I \mapsto I \cap Q)$, $\mathcal{D}_K(H, Q)$ is equal to $\mathcal{D}_K(\varphi)$. In particular, $\mathcal{D}_K(H, Q)$ is the ordinal doset Hibi ring.

Here we consider a generating system of $\mathcal{D}_K(H, Q)$ as a K -subalgebra of $\mathcal{R}_K(H)$. Set $H_1 := \{\alpha \in H \mid y \leq \alpha \text{ for some } y \in Q\}$, $H_2 = H \setminus H_1$ and $\psi: H_1 \rightarrow J(Q) \setminus \{\emptyset\}$, $(\alpha \mapsto \{y \in Q \mid y \leq \alpha\})$.

Lemma 4.11 *Let $\alpha_1 \alpha_2 \cdots \alpha_r$ be a standard monomial on H with $\alpha_1 \leq \alpha_2 \leq \cdots \leq \alpha_r$, $\alpha_1, \dots, \alpha_s \in H_2$ and $\alpha_{s+1}, \dots, \alpha_r \in H_1$. Then $\alpha_1 \cdots \alpha_r \in \mathcal{D}_K(H, Q)$ if and only if $r - s \equiv 0 \pmod{2}$ and $\psi(\alpha_{s+2i-1}) = \psi(\alpha_{s+2i})$ for $i = 1, 2, \dots, (r - s)/2$.*

Proof Let ν be the element of $\overline{\mathcal{T}}(P)$ corresponding to $\alpha_1 \cdots \alpha_r$. Then $\nu(x) = \#\{j \mid x \leq \alpha_j\}$. So $\nu(y) = \#\{j \mid j > s \text{ and } \psi(\alpha_j) \ni y\}$ for $y \in Q$. Since $\emptyset \neq \psi(\alpha_{s+1}) \subset \psi(\alpha_{s+2}) \subset \cdots \subset \psi(\alpha_r)$, $\nu(y) \equiv 0 \pmod{2}$ for any $y \in Q$ if and only if $r - s \equiv 0 \pmod{2}$ and $\psi(\alpha_{s+2i-1}) = \psi(\alpha_{s+2i})$ for $i = 1, 2, \dots, (r - s)/2$. The result follows from Theorem 4.10. ■

Now assume that Q has a unique minimal element y_0 . Then $H_1 = \{\alpha \in H \mid \alpha \geq y_0\}$ and $H_2 = \{\alpha \in H \mid \alpha \not\geq y_0\}$.

Lemma 4.12 (1) *If $\alpha, \beta \in H_1$, then $\alpha \wedge \beta, \alpha \vee \beta \in H_1$.*

(2) *If $\alpha \in H_1$ and $\beta \in H_2$, then $\alpha \wedge \beta \in H_2$, $\alpha \vee \beta \in H_1$ and $\psi(\alpha \vee \beta) = \psi(\alpha)$.*

(3) *If $\alpha, \beta \in H_2$, then $\alpha \wedge \beta, \alpha \vee \beta \in H_2$.*

(4) *If $\alpha, \beta \in H_1$, then $\psi(\alpha \wedge \beta) = \psi(\alpha) \cap \psi(\beta)$ and $\psi(\alpha \vee \beta) = \psi(\alpha) \cup \psi(\beta)$.*

(5) *If $\alpha, \beta, \alpha', \beta' \in H_1$, $\psi(\alpha) = \psi(\beta)$ and $\psi(\alpha') = \psi(\beta')$, then $\psi(\alpha \wedge \alpha') = \psi((\alpha \vee \alpha') \wedge (\beta \wedge \beta'))$ and $\psi(\beta \vee \beta') = \psi((\alpha \vee \alpha') \vee (\beta \wedge \beta'))$.*

Proof Let δ be an element of H and I the corresponding element of $J(P) \setminus \{\emptyset\}$ by the lattice isomorphism $H \simeq J(P) \setminus \{\emptyset\}$. Then $\delta \in H_1$ if and only if $I \ni y_0$, and $\wedge$ and $\vee$ operations in H correspond to $\cap$ and $\cup$ in $J(P) \setminus \{\emptyset\}$ respectively. Furthermore, $\psi(\delta) = I \cap Q$ if $\delta \in H_1$. The lemma is proved straightforwardly by these facts. ■

Now we state the following

Proposition 4.13 *If Q has a unique minimal element, then $\mathcal{D}_K(H, Q)$ is a K -subalgebra of $\mathcal{R}_K(H)$ generated by $\{\alpha\beta \mid \alpha, \beta \in H_1, \alpha \leq \beta \text{ and } \psi(\alpha) = \psi(\beta)\} \cup H_2$.*

Proof We set $G = \{\alpha\beta \mid \alpha, \beta \in H_1, \alpha \leq \beta \text{ and } \psi(\alpha) = \psi(\beta)\} \cup H_2$. The by Lemma 4.11, we see that $\mathcal{D}_K(H, Q) \subset K[G]$.

To prove the reverse inclusion, let μ be an arbitrary product of elements of G . Then by Lemma 4.12 and equation (4.1), we see that, by repeated application of the straightening law of $\mathcal{R}_K(H)$, μ is equal to a standard monomial $\alpha_1 \cdots \alpha_r$ with $\alpha_1 \leq \cdots \leq \alpha_r$ such that for some s , $\alpha_1, \dots, \alpha_s \in H_2$, $\alpha_{s+1}, \dots, \alpha_r \in H_1$, $r-s \equiv 0 \pmod{2}$ and $\psi(\alpha_{s+2i-1}) = \psi(\alpha_{s+2i})$ for $i = 1, 2, \dots, (r-s)/2$. Then again by Lemma 4.11, we see that $\alpha_1 \cdots \alpha_r \in \mathcal{D}_K(H, Q)$. Therefore $\mu \in \mathcal{D}_K(H, Q)$. ■

Now we apply the theory of (generalized) doset Hibi rings to the ring of $O(m)$ and $SO(m)$ invariants.

Set $L = \{1, 2, \dots, m\}$ with reverse order and $\varphi: \Gamma'(m \times n; \gamma) \rightarrow L$, $(\delta \mapsto \text{size}\delta)$. Then φ is a lattice homomorphism and $\varphi^*(j) = [b_1, b_2, \dots, b_j]$. By analyzing the relations of the leading monomials of the sagbi basis $\{[\alpha|\beta]_{Z_\gamma^T Z_\gamma} \mid \alpha, \beta \in \Gamma'(m \times n; \gamma), \text{size}\alpha = \text{size}\beta\}$ of $K[Z_\gamma]^{O(m, -)} = K[Z_\gamma^T Z_\gamma]$, and the sagbi basis $\{[\alpha|\beta]_{Z_\gamma^T Z_\gamma} \mid \alpha, \beta \in \Gamma'(m \times n; \gamma), \text{size}\alpha = \text{size}\beta < m\} \cup \{\delta_{Z_\gamma} \mid \delta \in \Gamma(m \times n; \gamma)\}$ of $K[Z_\gamma]^{SO(m, -)} = K[Z_\gamma^T Z_\gamma, \Gamma(Z_\gamma)]$, we see the following

Theorem 4.14 (1) *in $K[Z_\gamma]^{O(m, -)} = \text{in } K[Z_\gamma^T Z_\gamma]$ is the doset Hibi ring over K defined by φ .*

(2) *in $K[Z_\gamma]^{SO(m, -)} = \text{in } K[Z_\gamma^T Z_\gamma, \Gamma(Z_\gamma)]$ is the generalized doset Hibi ring defined by $\Gamma'(m \times n; \gamma)$ and $\{[b_1, \dots, b_{m-1}], \dots, [b_1, b_2], [b_1]\}$.*

In particular, in $K[Z_\gamma]^{O(m, -)}$ and in $K[Z_\gamma]^{SO(m, -)}$ are normal affine semi-group rings and therefore $K[Z_\gamma]^{O(m, -)}$ and $K[Z_\gamma]^{SO(m, -)}$ are normal and Cohen-Macaulay, and has rational singularities if $\text{char}K = 0$ and is F -rational if $\text{char}K > 0$.

5 Gorenstein property

In this section, we stat a criterion of the Gorenstein property of a (generalized) doset Hibi ring.

Let H be a finite distributive lattice, P the set of join-irreducible elements of H and Q a subset of P and $\{T_x\}_{x \in P}$ the family of indeterminates indexed by P . Set $\overline{\mathcal{T}}(P, Q) := \{\nu \in \overline{\mathcal{T}}(P) \mid \nu(y) \equiv 0 \pmod{2} \text{ for any } y \in Q\}$ and $\mathcal{T}(P, Q) := \{\nu \in \mathcal{T}(P) \mid \nu(y) \equiv 0 \pmod{2} \text{ for any } y \in Q\}$. Then since $\mathcal{D}_K(H, Q) = \bigoplus_{\nu \in \overline{\mathcal{T}}(P, Q)} K T^\nu$, we see, by Theorem 2.2, that the canonical module of $\mathcal{D}_K(H, Q)$ is $\bigoplus_{\nu \in \mathcal{T}(P, Q)} K T^\nu$.

We define a order on $\mathcal{T}(P, Q)$ by $\nu \leq \nu' \stackrel{\text{def}}{\iff} \nu' - \nu \in \overline{\mathcal{T}}(P, Q)$, where $(\nu' - \nu)(x) := \nu'(x) - \nu(x)$ for any $x \in P$. Then it is clear that T^ν is a generator of the canonical module of $\mathcal{D}_K(H, Q)$ if and only if ν is a minimal element of $\mathcal{T}(P, Q)$. In particular, $\mathcal{D}_K(H, Q)$ is Gorenstein if and only if $\mathcal{T}(P, Q)$ has a unique minimal element.

Now we set $P^+ := P \cup \{\infty\}$, where ∞ is a new element and $x < \infty$ for any $x \in P$ and $Q^+ := Q \cup \{\infty\}$. We also set $\tilde{P} := P^+ \cup \{(y_1, y_2) \in Q^+ \times Q^+ \mid y_1 < y_2 \text{ in } P^+\}$ and define the order on $\tilde{P}$ by extending the relation $y_1 < (y_1, y_2) < y_2$ for $y_1, y_2 \in Q^+$ with $y_1 < y_2$ in P^+ . Then we have the following result.

Theorem 5.1 $\mathcal{D}_K(H, Q)$ is Gorenstein if and only if the following two conditions are satisfied.

- (1) $\tilde{P}$ is pure.
- (2) For any $y, y' \in Q^+$ with $y < y'$, $\text{rank}[y, y']_{\tilde{P}} \equiv 0 \pmod{2}$.

Proof First we define $\tilde{\nu}_0: \tilde{P} \rightarrow \mathbf{N}$ by downward induction as follows.

- $\tilde{\nu}_0(\infty) = 0$.
- If $x \neq \infty$ and $\tilde{\nu}_0(x')$ is defined for any $x' \in \tilde{P}$ with $x' > x$, we set

$$\tilde{\nu}_0(x) = \begin{cases} \max\{\tilde{\nu}_0(x') + 1 \mid x' > x\} & \text{if } x \notin Q, \\ 2\lceil \max\{\tilde{\nu}_0(x') + 1 \mid x' > x\} / 2 \rceil & \text{if } x \in Q. \end{cases}$$

Then it is clear that $\nu_0 := \tilde{\nu}_0|_P \in \mathcal{T}(P, Q)$ and $\nu(x) \geq \nu_0(x)$ for any $\nu \in \mathcal{T}(P, Q)$ and $x \in P$. In particular, ν_0 is a minimal element of $\mathcal{T}(P, Q)$.

Now we start the proof of the theorem and we first prove the “if” part of the theorem. By assumptions (1) and (2), it is easily verified that $\tilde{\nu}_0(x) = \tilde{\nu}_0(x') + 1$ for any $x, x' \in \tilde{P}$ with $x < x'$. Therefore, by the definition of $\tilde{P}$, we see that $\nu - \nu_0 \in \overline{\mathcal{T}}(P, Q)$ for any $\nu \in \mathcal{T}(P, Q)$. So ν_0 is the unique minimal element of $\mathcal{T}(P, Q)$ and $\mathcal{D}_K(H, Q)$ is Gorenstein.

Next we prove the “only if” part. If (1) or (2) is not valid, then there are $x, x' \in \tilde{P}$ with $x < x'$ in $\tilde{P}$ such that $\tilde{\nu}_0(x) \geq \tilde{\nu}_0(x') + 2$. Since $x' \neq \infty$ in such a case, we see that there are $x, x' \in P$ such that $x < x'$ in P , $\{x, x'\} \not\subset Q$ and $\nu_0(x) \geq \nu_0(x') + 2$ or there are $y, y' \in Q$ such that $y < y'$ in P and $\nu_0(y) \geq \nu_0(y') + 4$. We classify this phenomena into three cases.

Case 5.1.1 There are $y, y' \in Q$ such that $y < y'$ in P and $\nu_0(y) \geq \nu_0(y') + 4$.

Case 5.1.2 There are $x, x' \in P$ such that $x < x'$, $x \notin Q$ and $\nu_0(x) \geq \nu_0(x') + 2$.

Case 5.1.3 There are $x, x' \in P$ such that $x < x'$, $x \in Q$, $x' \notin Q$ and $\nu_0(x) \geq \nu_0(x') + 2$.

As for Case 5.1.1, we set

$$\nu(z) = \begin{cases} \nu_0(z) & z \not\leq y' \text{ or } z = y, \\ \nu_0(z) + 2 & z \leq y' \text{ and } z \neq y. \end{cases}$$

Then it is easily verified that $\nu \in \mathcal{T}(P, Q)$ and $\nu - \nu_0 \notin \overline{\mathcal{T}}(P, Q)$ since $(\nu - \nu_0)(y) = (\nu - \nu_0)(y') - 2$. Therefore, there is a minimal element of $\mathcal{T}(P, Q)$ other than ν_0 .

As for Case 5.1.2, we set

$$\nu(z) = \begin{cases} \nu_0(z) & z \not\leq x', \\ \nu_0(z) + 1 & z = x, \\ \nu_0(z) + 2 & z \leq x' \text{ and } z \neq x. \end{cases}$$

Then it is easily verified that $\nu \in \mathcal{T}(P, Q)$ and $\nu - \nu_0 \notin \overline{\mathcal{T}}(P, Q)$ since $(\nu - \nu_0)(x) = (\nu - \nu_0)(x') - 1$. Therefore, there is a minimal element of $\mathcal{T}(P, Q)$ other than ν_0 .


Finally, as for Case 5.1.3, we set

$$\nu(z) = \begin{cases} \nu_0(z) & z \not\leq x' \text{ or } z = x, \\ \nu_0(z) + 1 & z = x', \\ \nu_0(z) + 2 & z < x' \text{ and } z \neq x. \end{cases}$$


Again it is easily verified that $\nu \in \mathcal{T}(P, Q)$ and $\nu - \nu_0 \notin \overline{\mathcal{T}}(P, Q)$. Therefore, there is a minimal element of $\mathcal{T}(P, Q)$ other than ν_0 .

In any case, we see that $\mathcal{D}_K(H, Q)$ is not Gorenstein. ■

Example 5.2 (1) If




where big dots express elements of Q , then




Therefore, $\mathcal{D}_K(H, Q)$ is Gorenstein.

(2) If




then




Therefore, $\tilde{P}$ is not pure and $\mathcal{D}_K(H, Q)$ is not Gorenstein.

(3) If



then



So there are $y, y' \in Q^+$ such that $y < y'$ and $\text{rank}[y, y']_{\tilde{P}} = 3$. Therefore, $\mathcal{D}_K(H, Q)$ is not Gorenstein.

Finally, we apply Theorem 5.1 to obtain a criterion of Gorenstein property of in $K[Z_\gamma]^{\text{O}(m, -)}$ and in $K[Z_\gamma]^{\text{SO}(m, -)}$. Note by Corollary 2.4, $K[Z_\gamma]^{\text{O}(m, -)}$ (or $K[Z_\gamma]^{\text{SO}(m, -)}$ resp.) is Gorenstein if and only if so is in $K[Z_\gamma]^{\text{O}(m, -)}$ (or in $K[Z_\gamma]^{\text{SO}(m, -)}$ resp.).

First we note the following lemma which is easily proved.

Lemma 5.3 *Let H be a distributive lattice and x an element of H which is not minimal. Then x is join-irreducible if and only if there uniquely exists an element $\alpha \in H$ such that $\alpha \lessdot x$.*

In order to apply Theorem 5.1, we first analyze the structure of the set of join-irreducible elements of $\Gamma'(m \times n; \gamma)$. Let P be the set of join-irreducible elements of $\Gamma'(m \times n; \gamma)$ and $\delta = [d_1, d_2, \dots, d_t]$ be an element of $\Gamma'(m \times n; \gamma) \setminus \{\gamma\}$.

First consider the case where $\text{size}\delta = m$. Then $[c_1, c_2, \dots, c_m] < [d_1, d_2, \dots, d_m]$ if and only if there is i such that $c_i = d_i - 1$ and $c_j = d_j$ for $j \neq i$. Therefore, by Lemma 5.3, we see that $\delta = [d_1, \dots, d_m]$ is join-irreducible if and only if there is a unique i such that

$$d_i > b_i \quad \text{and} \quad d_i > d_{i-1} + 1,$$

where we set $d_0 := 0$ (recall that $\gamma = [b_1, \dots, b_m]$). Set $\xi(\delta) := (n - d_i - m + i, i - 1)$ for such δ . Note $n - d_i - m + i = \#\{j \mid j \notin \{d_1, \dots, d_m\} \text{ and } d_i < j \leq n\}$ and $i - 1 = \#\{j \mid j \in \{d_1, \dots, d_m\} \text{ and } 1 \leq j < d_i\}$. Then it is easily verified that ξ is a poset anti-homomorphism from $\{\delta \in P \setminus \{\gamma\} \mid \text{size}\delta = m\}$ to $\mathbf{N} \times \mathbf{N}$.

Next consider the case where $\text{size}\delta < m$. Then $[c_1, \dots, c_{t'}] < [d_1, \dots, d_t]$ if and only if

- (1) $t' = t + 1$, $c_{t'} = m$ and $c_i = d_i$ for $i = 1, 2, \dots, t$ or
- (2) $t' = t$ and there is i such that $c_i = d_i - 1$ and $c_j = d_j$ for $j \neq i$.

Therefore, δ is join-irreducible if and only if $\delta = [b_1, b_2, \dots, b_t]$ or there is i such that

$$d_j = b_j \text{ for } j < i \text{ and } d_j = n - t + j \text{ for } j \geq i.$$

We extend ξ by setting $\xi(\delta) := (t - m, i - 1)$ for the latter case and $\xi([b_1, \dots, b_t]) := (t - m, t)$. Then it is easily verified that ξ is a poset anti-homomorphism from $P \setminus \{\gamma\}$ to $\mathbf{Z} \times \mathbf{N}$.


Now we separate $\{b_1, b_2, \dots, b_m\}$ into blocks and define the gaps between them. Set $\{u \mid 1 \leq u \leq m, b_u + 1 < b_{u+1}\} = \{u(1), \dots, u(k)\}$ with $u(1) < \dots < u(k)$, where we set $b_{m+1} := n + 1$, and $\chi_0 := \{1, 2, \dots, b_1 - 1\}$, $B_1 := \{b_1, b_2, \dots, b_{u(1)}\}$, $\chi_1 := \{b_{u(1)} + 1, b_{u(1)} + 2, \dots, b_{u(1)+1} - 1\}$, $B_2 := \{b_{u(1)+1}, b_{u(1)+2}, \dots, b_{u(2)}\}$, $\chi_2 := \{b_{u(2)} + 1, b_{u(2)} + 2, \dots, b_{u(2)+1} - 1\}$, $B_3 := \{b_{u(2)+1}, b_{u(2)+2}, \dots, b_{u(3)}\}$, $\dots$, $B_k := \{b_{u(k-1)+1}, b_{u(k-1)+2}, \dots, b_{u(k)}\}$, $\chi_k := \{b_{u(k)} + 1, b_{u(k)} + 2, \dots, b_{u(k)+1} - 1\}$ and $B_{k+1} := \{b_{u(k)+1}, b_{u(k)+2}, \dots, b_m\}$. Note that if $b_m < n$, then $u(k) = m$ and $B_{k+1} = \emptyset$.

Example 5.4 (1) If $m = 7$, $n = 15$ and $\gamma = [2, 6, 7, 8, 10, 13, 14]$, then $k = 4$, $u(1) = 1$, $u(2) = 4$, $u(3) = 5$, $u(4) = 7$ and $\chi_0 = \{1\}$, $B_1 = \{2\}$, $\chi_1 = \{3, 4, 5\}$, $B_2 = \{6, 7, 8\}$, $\chi_2 = \{9\}$, $B_3 = \{10\}$, $\chi_3 = \{11, 12\}$, $B_4 = \{13, 14\}$, $\chi_4 = \{15\}$ and $B_5 = \emptyset$.


- (2) If $m = 7$, $n = 14$ and $\gamma = [2, 6, 7, 8, 10, 13, 14]$, then $k = 3$, $u(1) = 1$, $u(2) = 4$, $u(3) = 5$ and $\chi_0 = \{1\}$, $B_1 = \{2\}$, $\chi_1 = \{3, 4, 5\}$, $B_2 = \{6, 7, 8\}$, $\chi_2 = \{9\}$, $B_3 = \{10\}$, $\chi_3 = \{11, 12\}$ and $B_4 = \{13, 14\}$,

It is easy to see that minimal elements of $P \setminus \{\gamma\}$ are $[b_1, \dots, b_{u(1)-1}, b_{u(1)} + 1, b_{u(1)+1}, \dots, b_m]$, $[b_1, \dots, b_{u(2)-1}, b_{u(2)} + 1, b_{u(2)+1}, \dots, b_m]$, $\dots$, $[b_1, \dots, b_{u(k-1)-1}, b_{u(k-1)} + 1, b_{u(k-1)+1}, \dots, b_m]$ and $[b_1, \dots, b_{m-1}, b_m + 1]$ in the case where $b_m < n$ and $[b_1, \dots, b_{u(1)-1}, b_{u(1)} + 1, b_{u(1)+1}, \dots, b_m]$, $[b_1, \dots, b_{u(2)-1}, b_{u(2)} + 1, b_{u(2)+1}, \dots, b_m]$, $\dots$, $[b_1, \dots, b_{u(k)-1}, b_{u(k)} + 1, b_{u(k)+1}, \dots, b_m]$ and $[b_1, b_2, \dots, b_{m-1}]$ in the case where $b_m = n$. Therefore, by considering the image of ξ , we see the Hasse diagram of $P \setminus \{\gamma\}$ is as indicated by the following example.

Example 5.5 (1) If $m = 7$, $n = 15$ and $\gamma = [2, 6, 7, 8, 10, 13, 14]$, then the Hasse diagram of $P \setminus \{\gamma\}$ is the following.



- (2) If $m = 7$, $n = 14$ and $\gamma = [2, 6, 7, 8, 10, 13, 14]$, then the Hasse diagram of $P \setminus \{\gamma\}$ is the following.



Set $Q_1 := \{[b_1, \dots, b_i] \mid i = 1, \dots, m\}$ and $Q_2 := \{[b_1, \dots, b_i] \mid i = 1, \dots, m-1\}$. Then by Theorem 4.14, we see that in $K[Z_\gamma]^{\text{O}(m,-)} \simeq D_K(\Gamma'(m \times n; \gamma), Q_1)$ and in $K[Z_\gamma]^{\text{SO}(m,-)} \simeq D_K(\Gamma'(m \times n; \gamma), Q_2)$. So by Theorem 5.1 and Corollary 2.4, we see the following results.

Theorem 5.6 ([Con, Corollary 2.3]) (1) If $b_m < n$, then $K[Z_\gamma]^{\text{O}(m,-)} = K[Z_\gamma^T Z_\gamma]$ is Gorenstein if and only if $|B_i| = |\chi_{i-1}|$ for $i = 2, 3, \dots, k$ and $|\chi_k| \equiv 1 \pmod{2}$.

(2) If $b_m = n$, then $K[Z_\gamma]^{\text{O}(m,-)} = K[Z_\gamma^T Z_\gamma]$ is Gorenstein if and only if $|B_i| = |\chi_{i-1}|$ for $i = 2, 3, \dots, k$ and $|B_{k+1}| = |\chi_k| - 1$.

Theorem 5.7 (1) If $b_m < n$, then $K[Z_\gamma]^{\text{SO}(m,-)} = K[Z_\gamma^T Z_\gamma, \Gamma(Z_\gamma)]$ is Gorenstein if and only if $|B_i| = |\chi_{i-1}|$ for $i = 2, 3, \dots, k$.

(2) If $b_m = n$, then $K[Z_\gamma]^{\text{SO}(m,-)} = K[Z_\gamma^T Z_\gamma, \Gamma(Z_\gamma)]$ is Gorenstein if and only if $|B_i| = |\chi_{i-1}|$ for $i = 2, 3, \dots, k+1$.

Let $\tilde{P}_i$ be the poset defined as in the preceding paragraph of Theorem 5.1 by P and Q_i for $i = 1, 2$. Then, the case where $b_m < n$, $\tilde{P}_1 = \tilde{P}_2 = P \cup \{\infty\}$. So the last condition of (1) in Theorem 5.6 corresponds to the condition (2) of Theorem 5.1 for $[\gamma, [b_1, \dots, b_{m-1}]]_{\tilde{P}_1}$. As for the case $b_m = n$, $\tilde{P}_1$ is obtained by adding a new element to $\tilde{P}_2$. So there is a difference between the condition for pureness of $\tilde{P}_1$ and $\tilde{P}_2$. Note that the condition in (2) of Theorem 5.7 can be written as $|B_i| = |\chi_{i-1}|$ for $i = 2, 3, \dots, k$ and $|B_{k+1}| = |\chi_k|$.

References

- [BC] Bruns, W. and Conca, A.: *F-rationality of determinantal rings and their Rees rings*. Michigan Math. J. **45** (1998) 291–299.
- [BH] Bruns, W. and Herzog, J.: “Cohen-Macaulay rings.” Cambridge studies in advanced mathematics **39** Cambridge University Press (1993)
- [BV] Bruns, W. and Vetter, U.: “Determinantal Rings.” Lecture Notes in Mathematics **1327** Springer (1988)
- [BRW] Bruns, W. Römer, T. and Wiebe, A: *Initial algebras of determinantal rings, Cohen-Macaulay and Ulrich ideals*. Michigan Math. J. **53** (2005), 71–81.
- [Con] Conca, A: *Divisor class group and the canonical class of determinantal rings defined by ideals of minors of a symmetric matrix* Arch. Math. **63** (1994), 216–224.
- [CHV] Conca, A., Herzog, J. and Valla, G.: *Sagbi bases with applications to blow-up algebras*. J. reine angew. Math. **474** (1996), 113–138.
- [DEP1] DeConcini, C., Eisenbud, D. and Procesi, C.: *Young Diagrams and Determinantal Varieties*. Inv. Math. **56** (1980), 129–165
- [DEP2] DeConcini, C., Eisenbud, D. and Procesi, C.: “Hodge Algebras.” Astérisque **91** (1982)
- [DP] DeConcini, C. and Procesi, C.: *A characteristic-free approach to invariant theory*, Adv. Math. **21** (1976), 330–354.
- [Hib] Hibi, T.: *Distributive lattices, affine smigroup rings and algebras with straightening laws*. in “Commutative Algebra and Combinatorics” (M. Nagata and H. Matsumura, ed.), Advanced Studies in Pure Math. **11** North-Holland, Amsterdam (1987), 93–109.
- [Hoc] Hochster, M.: *Rings of invariants of tori, Cohen-Macaulay rings generated by monomials and polytopes*. Ann. of Math. **96** (1972), 318–337
- [Miy] Miyazaki, M: *A Note on the Invariants of the Group of Triangular Matrices Acting on Generic Echelon Matrices*, Bulletin of Kyoto University of Education Ser. B **85** (1994), 57–62.

- [Sta] Stanley, R. P.: *Hilbert Functions of Graded Algebras*. Adv. Math.
28 (1978), 57–83.